

A STUDY ON FUZZY ROUGH ALGEBRAIC BORDER AND EXTERIOR

M. SRIVIDYA and G. JAYALALITHA

Assistant Professor Sona College of Arts and Science Salem, Tamil Nadu, India E-mail: srividyamohan38@gmail.com

Professor, Vels Institute of Science Technology and Advanced Studies (VISTAS) Chennai, Tamil Nadu, India E-mail: g.jayalalithamaths.sbs@velsuniv.ac.in

Abstract

The concepts of fuzzy rough algebraic border and fuzzy rough algebraic exterior sets are introduced in this paper. The properties of fuzzy rough algebraic border and fuzzy rough algebraic exterior are discussed. There are some intriguing characterizations and attributes studied.

1. Introduction

Authors [14], [8], [9], and [3] discussed fuzzy topological spaces, fuzzy sets, and their applications. Soft set theory, intuitionistic fuzzy set theory, soft fuzzy set theory, and other approaches are used to address a variety of additional uncertainties that crop up in real-world situations. In [12] and [13], a few of such theories' characteristics are covered. The idea of rough set was first presented by Pawlak [7]. R. Biswas and S. Nanda investigated the rough group and rough subgroup [1]. The rough topological space was researched by B.P. Mathew and S. J. John [5]. S. Majmudar and S. Nanda [6] examined the idea of a fuzzy rough set. The fuzzy rough topological group and

2020 Mathematics Subject Classification: 54A40, 03E72.

Keywords: fuzzy rough algebraic border, fuzzy rough algebraic exterior, fuzzy rough algebraic boundary.

Received January 5, 2024; Accepted January 16, 2024

M. SRIVIDYA and G. JAYALALITHA

fuzzy rough group were examined and discussed by the author [11]. Calda, Jafari, and Noiri [2] researched the ideas of G-exterior and G-border. The concepts of fuzzy rough algebraic border and fuzzy rough algebraic exterior sets are introduced in this paper. The properties of fuzzy rough algebraic border and fuzzy rough algebraic exterior are discussed. There are some intriguing characterizations and attributes studied.

2. Preliminaries

Definition 2.1 [11]. Let X be the fuzzy topological space, and let be a fuzzy set. The boundary Bd of λ , is then determined as $Bd(\lambda) = cl(\lambda) \cap cl(\lambda)'$. Obviously, $Bd(\lambda)$ is a fuzzy closed set.

Definition 2.2 [2]. The fuzzy rough border of any fuzzy rough set in (X, T) is defined as the intersection of the fuzzy rough set with the closure of its compliment.

Definition 2.3 [2]. The fuzzy rough exterior of any fuzzy rough set in (X, T) is defined as the interior of the compliment of the fuzzy rough set.

3. Fuzzy Rough Algebraic Border and Exterior

Definition 3.1. Let (X, TM) be a fuzzy rough algebraic TM system and be any fuzzy rough algebraic. Then the fuzzy rough algebraic boundary of A, is denoted and defined as

$$\mathcal{F}_{\mathcal{RTM}}Bd(A) = \mathcal{F}_{\mathcal{RTM}}cl(A) \cap \mathcal{F}_{\mathcal{RTM}}cl(A).$$

Definition 3.2. Let A be a fuzzy rough algebraic in a fuzzy rough algebraic TM system (X, TM). Then the fuzzy rough algebraic border of A is defined and denoted by

$$\mathcal{F}_{\mathcal{RTM}}Br(A) = A \cap \mathcal{F}_{\mathcal{RTM}}cl(A).$$

Proposition 3.1. Let A be any fuzzy rough algebraic in a fuzzy rough algebraic TM system (X, TM). Then

(i)
$$\mathcal{F}_{\mathcal{RTM}}Br(A) \subset \mathcal{F}_{\mathcal{RTM}}cl(A)'.$$

(ii)
$$\mathcal{F}_{\mathcal{RTM}} \operatorname{int}(\mathcal{F}_{\mathcal{RTM}}Br(A)) \subseteq A$$
.

(iii)
$$\mathcal{F}_{\mathcal{RTM}}Br(\mathcal{F}_{\mathcal{RTM}}Br(A)) \subseteq \mathcal{F}_{\mathcal{RTM}}Br(A).$$

Proof.

i)
$$\mathcal{F}_{\mathcal{RTM}}Br(A) = A \cap \mathcal{F}_{\mathcal{RTM}}cl(A)'$$

 $\subseteq \mathcal{F}_{\mathcal{RTM}} cl(A')$

Hence
$$\mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq \mathcal{F}_{\mathcal{RTM}}cl(A')$$
.

```
ii) \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(\mathcal{F}_{\mathcal{RTM}} Bd(A)) = \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A \cap \mathcal{F}_{\mathcal{RTM}} cl(A'))
```

 $\subseteq A \cap \mathcal{F}_{\mathcal{RTM}}cl(A')$

```
= A
```

Hence $\mathcal{F}_{\mathcal{RTM}} \operatorname{int}(\mathcal{F}_{\mathcal{RTM}}Br(A)) \subseteq A$.

iii)
$$\mathcal{F}_{\mathcal{RTM}}Br(\mathcal{F}_{\mathcal{RTM}}Br(A)) = \mathcal{F}_{\mathcal{RTM}}Br(A \cap \mathcal{F}_{\mathcal{RTM}}cl(A'))$$

 $\subseteq (A \cap \mathcal{F}_{\mathcal{RTM}} cl(A')) \cap (\mathcal{F}_{\mathcal{RTM}} cl(A \cap \mathcal{F}_{\mathcal{RTM}} cl(A'))')$

 $\subseteq A \cap \mathcal{F}_{\mathcal{RTM}} cl(A')$

Hence
$$\mathcal{F}_{\mathcal{RTM}}Br(\mathcal{F}_{\mathcal{RTM}}Br(A)) \subseteq \mathcal{F}_{\mathcal{RTM}}Br(A)$$
.

Proposition 3.2. If A is any fuzzy rough algebraic open in (X, TM) then $\mathcal{F}_{RTM}Br(A) \subseteq A$.

Proof. Since A is a fuzzy rough algebraic open then A' is a fuzzy rough algebraic closed. Then

$$\mathcal{F}_{\mathcal{RTM}}Br(A) = A \cap \mathcal{F}_{\mathcal{RTM}}cl(A')$$

 $\subseteq A \cap A'$
 $\subseteq A.$

Hence $\mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq A$.

Proposition 3.3. Let C and D be any two fuzzy rough algebraic in (X, TM) then $\mathcal{F}_{\mathcal{RTM}}Br(C) \cup \mathcal{F}_{\mathcal{RTM}}Br(D)$.

Proof.

$$\begin{split} \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(C \cup D) &= (C \cup D) \cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(C \cup D)^{'} \\ &= (C \cup D) \cap (\mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(C^{'} \cap D^{'})) \\ &\subseteq (C \cup D) \cap (\mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(C^{'}) \cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(D^{'})) \\ &= (\mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(C^{'}) \cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(D^{'})) \\ &\cup (\mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(D) \cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(C^{'})) \\ &= \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(C) \cup \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(D) \end{split}$$

Hence $\mathcal{F}_{\mathcal{RTM}}Br(C \cup D) \subseteq \mathcal{F}_{\mathcal{RTM}}Br(C') \cup \mathcal{F}_{\mathcal{RTM}}Br(D').$

Proposition 3.4. Let S and T be any two fuzzy rough algebraic in fuzzy rough TM system (X, TM). Then

$$\mathcal{F}_{\mathcal{RTM}}Br(S\cap T) \supseteq \mathcal{F}_{\mathcal{RTM}}Br(S) \cap \mathcal{F}_{\mathcal{RTM}}Br(T).$$

Proof.

$$\begin{split} \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(S\cap T) &= (S\cap T)\cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(S\cap T)^{'} \\ &= (S\cap T)\cap (\mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(S^{\prime}\cup T^{\prime})) \\ &= (S\cap T)\cap (\mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(S^{\prime})\cup \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(T^{\prime})) \\ &\supseteq (S\cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(S^{\prime}))\cap (T\cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(T^{\prime})) \\ &= \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(S)\cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(T). \end{split}$$

Hence $\mathcal{F}_{\mathcal{RTM}}Br(S \cap T) \supseteq \mathcal{F}_{\mathcal{RTM}}Br(S) \cap \mathcal{F}_{\mathcal{RTM}}Br(T)$.

Advances and Applications in Mathematical Sciences, Volume 23, Issue 8, June 2024

742

Proposition 3.5. For any fuzzy rough algebraic A in (X, TM),

$$\mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq \mathcal{F}_{\mathcal{RTM}}Bd(A).$$

Proof.

$$\begin{aligned} \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Br(A) &= A \cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(A') \\ &\subseteq \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(A) \cap \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}cl(A') \\ &= \mathcal{F}_{\mathcal{R}\mathcal{T}\mathcal{M}}Bd(A). \end{aligned}$$

Therefore, $\mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq \mathcal{F}_{\mathcal{RTM}}Bd(A)$.

Remark 3.1. If A is any fuzzy rough algebraic closed in (X, TM) then $\mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq \mathcal{F}_{\mathcal{RTM}}Bd(A)$.

Proof. Since A is a fuzzy rough algebraic closed, $\mathcal{F}_{\mathcal{RTM}}cl(A) = A$.

Now,
$$\mathcal{F}_{\mathcal{RTM}}Br(A) = A \cap \mathcal{F}_{\mathcal{RTM}}cl(A')$$

= $\mathcal{F}_{\mathcal{RTM}}cl(A) \cap \mathcal{F}_{\mathcal{RTM}}cl(A')$.
= $\mathcal{F}_{\mathcal{RTM}}Bd(A)$.

Therefore, $\mathcal{F}_{\mathcal{RTM}}Br(A) = \mathcal{F}_{\mathcal{RTM}}Bd(A)$.

Definition 3.3. A fuzzy rough algebraic D in (X, TM) is said to be fuzzy rough algebraic exterior of D is denoted and defined by

$$\mathcal{F}_{\mathcal{RTM}}Er(D) = \mathcal{F}_{\mathcal{RTM}}\operatorname{int}(D').$$

Proposition 3.6. Let D be a fuzzy rough algebraic in fuzzy rough algebraic TM system (X, TM). Then

- (i) $\mathcal{F}_{\mathcal{RTM}}Er(D) \subseteq (\mathcal{F}_{\mathcal{RTM}}cl(D))'$.
- (ii) $\mathcal{F}_{\mathcal{RTM}}Er(\mathcal{F}_{\mathcal{RTM}}Er(D)) = \mathcal{F}_{\mathcal{RTM}}\operatorname{int}(\mathcal{F}_{\mathcal{RTM}}cl(D)).$
- (iii) $\mathcal{F}_{\mathcal{RTM}}Er(\widetilde{1}) = \widetilde{0}.$
- (iv) $\mathcal{F}_{\mathcal{RTM}}Er(\widetilde{0}) = \widetilde{1}$.

(v) $\mathcal{F}_{\mathcal{RTM}} \operatorname{int}(D) \subseteq \mathcal{F}_{\mathcal{RTM}} Er(\mathcal{F}_{\mathcal{RTM}} cl(D)).$

Proof.

(i) The proof follows directly from the Definition 2.17.

(ii)
$$\mathcal{F}_{\mathcal{RTM}}Er(\mathcal{F}_{\mathcal{RTM}}Er(D)) = \mathcal{F}_{\mathcal{RTM}}Er(\mathcal{F}_{\mathcal{RTM}}Er(D'))$$
 [By Definition 2.17]

 $= \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(\mathcal{F}_{\mathcal{RTM}} \operatorname{int}(D'))'$ $= \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(\mathcal{F}_{\mathcal{RTM}} cl(D)).$

Hence, $\mathcal{F}_{\mathcal{RTM}}Er(\mathcal{F}_{\mathcal{RTM}}Er(D)) = \mathcal{F}_{\mathcal{RTM}}\operatorname{int}(\mathcal{F}_{\mathcal{RTM}}cl(D)).$

(iii) By (ii),

$$\mathcal{F}_{\mathcal{RTM}}Er(\widetilde{1}) = \mathcal{F}_{\mathcal{RTM}}Er(\widetilde{1}') = \widetilde{0}.$$

(iv) By (ii),

$$\mathcal{F}_{\mathcal{RTM}}Er(\widetilde{0}) = \mathcal{F}_{\mathcal{RTM}}Er(\widetilde{0}') = \widetilde{1}.$$

(v) Since $A \subseteq \mathcal{F}_{\mathcal{RTM}} cl(A)$.

$$\Rightarrow \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A) \subseteq \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(\mathcal{F}_{\mathcal{RTM}} cl(A))$$
$$\Rightarrow \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A) \subseteq \mathcal{F}_{\mathcal{RTM}} Er(\mathcal{F}_{\mathcal{RTM}} Er(A)) \text{ by (iii).}$$

Proposition 3.7. Let R and S be any two fuzzy rough algebraic in a fuzzy rough algebraic TM system (X, TM). Then the following conditions hold.

(i) If
$$R \subseteq S$$
 then $\mathcal{F}_{\mathcal{RTM}}Er(R) \supseteq \mathcal{F}_{\mathcal{RTM}}Er(S)$.

(ii)
$$\mathcal{F}_{\mathcal{RTM}}Er(R \cup S) = \mathcal{F}_{\mathcal{RTM}}Er(R) \cap \mathcal{F}_{\mathcal{RTM}}Er(S).$$

(iii)
$$\mathcal{F}_{\mathcal{RTM}}Er(R \cap S) = \mathcal{F}_{\mathcal{RTM}}Er(R) \cup \mathcal{F}_{\mathcal{RTM}}Er(S)$$

Proof.

(i) Since $R \subseteq S$, $\mathcal{F}_{\mathcal{RTM}} Er(R) \subseteq \mathcal{F}_{\mathcal{RTM}} Er(S)$. Hence the proof is obvious by Definition 2.17.

Advances and Applications in Mathematical Sciences, Volume 23, Issue 8, June 2024

744

(ii)
$$\mathcal{F}_{\mathcal{RTM}} Er(R \cup S) = \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(R \cup S)'.$$

$$= \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(R' \cup S')$$

$$= \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(R') \cap \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(S')$$

$$= \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(R) \cap \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(S).$$

Hence, $\mathcal{F}_{\mathcal{RTM}}Er(R \cup S) = \mathcal{F}_{\mathcal{RTM}}Er(R) \cap \mathcal{F}_{\mathcal{RTM}}Er(S)$.

(iii) The proof is similar to (ii).

Proposition 3.8. Let A be a fuzzy rough algebraic in a fuzzy rough algebraic TM system (X, TM). Then $\mathcal{F}_{\mathcal{RTM}}Er(A) = A'$ if and only if A is fuzzy rough algebraic closed.

Proof. If *A* is any fuzzy rough algebraic *TM* closed then,

$$A = \mathcal{F}_{\mathcal{RTM}} cl(A).$$

By Definition 2.17, $\mathcal{F}_{\mathcal{RTM}}Er(A) = \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A')$

$$= \left(\mathcal{F}_{\mathcal{RTM}}(A) \right)'$$

= A'

Hence, $\mathcal{F}_{\mathcal{RTM}}Er(A) = A'$.

Conversely, $\mathcal{F}_{\mathcal{RTM}}Er(A) = A'$

$$\Rightarrow \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A') = A'.$$

Hence A' is a fuzzy rough algebraic open. Therefore, A is a fuzzy rough algebraic closed.

Proposition 3.9. Let A be any fuzzy rough algebraic in a fuzzy rough algebraic TM system (X, TM). Then

$$(\mathcal{F}_{\mathcal{RTM}}Bd(A))$$
 = $\mathcal{F}_{\mathcal{RTM}}$ int $(A) \cup \mathcal{F}_{\mathcal{RTM}}Er(A)$.

Proof.

Since, $\mathcal{F}_{\mathcal{RTM}}Bd(A) = \mathcal{F}_{\mathcal{RTM}}cl(A) \cap \mathcal{F}_{\mathcal{RTM}}cl(A').$

Then,
$$(\mathcal{F}_{\mathcal{RTM}}Bd(A))' = (\mathcal{F}_{\mathcal{RTM}}cl(A) \cap \mathcal{F}_{\mathcal{RTM}}cl(A'))'$$

= $\mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A') \cup \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A)$
 $\mathcal{T} = \operatorname{int}(A) \cup \mathcal{T} = \operatorname{int}(A')$

$$=\mathcal{F}_{\mathcal{RTM}}\operatorname{int}(A)\cup\mathcal{F}_{\mathcal{RTM}}\operatorname{int}(A')$$

$$= \mathcal{F}_{\mathcal{RTM}} \operatorname{int}(A) \cup \mathcal{F}_{\mathcal{RTM}} Er(A).$$

Therefore, $(\mathcal{F}_{\mathcal{RTM}}Bd(A))' = \mathcal{F}_{\mathcal{RTM}}\operatorname{int}(A) \cup \mathcal{F}_{\mathcal{RTM}}Er(A).$

Proposition 3.10. Let A be a fuzzy rough algebraic in a fuzzy rough algebraic TM system (X, TM). Then $\mathcal{F}_{\mathcal{RTM}}Br(A) = (\mathcal{F}_{\mathcal{RTM}}Er(A'))'$.

Proof.

Since, $\mathcal{F}_{\mathcal{RTM}}Br(A) = A \cap \mathcal{F}_{\mathcal{RTM}}cl(A')$

Then $\mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq A$

$$\subseteq \mathcal{F}_{\mathcal{RTM}} cl(A)$$

$$= (\mathcal{F}_{\mathcal{RTM}}Er(A))'.$$

Hence $\mathcal{F}_{\mathcal{RTM}}Br(A) = (\mathcal{F}_{\mathcal{RTM}}Er(A))'$.

Remark 3.2. From Propositions 3.5 to 3.10,

 $\mathcal{F}_{\mathcal{RTM}}Br(A) \not\subseteq \mathcal{F}_{\mathcal{RTM}}Br(A) \subseteq \mathcal{F}_{\mathcal{RTM}}Bd(A).$

References

- R. Biswas and S. Nanda, Rough groups and rough subgroups, Bull. Polish Acad. Sci. Math. 42 (1994), 251-254.
- [2] M. Caldas, S. Jafari and T. Noiri, Notions via g open sets, Kochi. J. Math. 2 (2007).
- [3] C. L. Chang, Fuzzy topological spaces, J. Math. Anal. Appl. 24 (1968), 182-190.

Advances and Applications in Mathematical Sciences, Volume 23, Issue 8, June 2024

746

- [4] T. K. Mondal and S. K. Samanta, Intuitionistic fuzzy rough sets and rough intuitionistic fuzzy sets, The Journal of Fuzzy Mathematics 9 (2001), 561-582.
- [5] BP Mathew and SJ John, On rough topological spaces, International Journal of Mathematical Archive 3(9) (2012), 3413-3421.
- [6] S. Nanda and S. Majumdar, Fuzzy rough sets, Fuzzy Sets and Systems 45 (1992), 157-160.
- [7] Z. Pawlak, Rough sets, Internat. J. Inform. Comput. Sci. 11(5) (1982), 341-356.
- [8] P. Smets, The degree of belief in a fuzzy event, Information Sciences 25 (1981), 1-19.
- [9] M. Sugeno, An introductory survey of fuzzy control, Information Sciences 36 (1985), 59-83.
- [10] D. Vidhya, E. Roja and MK Uma, Algebraic fuzzy rough sheaf group formed by pointed fuzzy rough topological group, Int. J. Math. and Comp. Appl. Research 4(1) (2014), 51-58.
- [11] D. Vidhya, E. Roja and MK Uma, On fuzzy rough BG boundary spaces, Annals of Fuzzy Mathematics and Informatics 10(4) (2015), 137-140.
- [12] T. Yogalakshmi, On pairwise intuitionistic fuzzy b disconnectedness, Global Journal of Pure and Applied Mathematics 12(3) (2016), 895-899.
- [13] T. Yogalakshmi, Disconnectedness in soft fuzzy centred systems, International Journal of Pure and Applied Mathematics 115(9) (2017), 223-229.
- [14] L. A. Zadeh, Fuzzy sets, Information and Control 8 (1965), 338-353.